

## Testing for Convergence or Divergence of a Series

Many of the series you come across will fall into one of several basic types. Recognizing these types will help you decide which tests or strategies will be most useful in finding whether a series is convergent or divergent.

### p-Series

$$\sum_{n=1}^{\infty} \frac{1}{n^p} \text{ is...}$$

- convergent if  $p > 1$
- divergent if  $p \leq 1$

### Geometric Series

$$\sum_{n=1}^{\infty} ar^{n-1} \text{ is...}$$

- convergent if  $|r| < 1$
- divergent if  $|r| \geq 1$

If  $a_n$  has a form that is similar to one of the above, see whether you can use the *comparison test*:

### Comparison Test

(Warning! This only works if  $a_n$  and  $b_n$  are always positive.)

- (i) If  $a_n \leq b_n$  for all  $n$ , and  $\sum b_n$  is convergent, then  $\sum a_n$  is convergent.  
 (ii) If  $a_n \geq b_n$  for all  $n$ , and  $\sum b_n$  is divergent, then  $\sum a_n$  is divergent.

**Example:**  $\sum_{n=1}^{\infty} \frac{1}{n^2 + n}$

Pick  $b_n = \frac{1}{n^2}$  (p-series)

$$a_n = \frac{1}{n^2 + n} \leq \frac{1}{n^2}, \text{ and}$$

$\sum_{n=1}^{\infty} \frac{1}{n^2}$  converges, so by

(i),  $\sum_{n=1}^{\infty} \frac{1}{n^2 + n}$  converges.

**Example:**  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$

Pick  $b_n = \frac{1}{2^n}$  (geometric)

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{2^n - 1} \frac{2^n}{1}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{1 - 1/2^n} = 1 > 0$$

$\sum_{n=1}^{\infty} \frac{1}{2^n}$  converges, so

$\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  converges.

Consider a series  $\sum b_n$  so that the ratio  $a_n/b_n$  cancels the dominant terms in the numerator and denominator of  $a_n$ , as in the example to the left. If you know whether  $\sum b_n$  converges or not, try using the limit comparison test.

### Limit Comparison Test

(Warning! This only works if  $a_n$  and  $b_n$  are always positive.)

If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c > 0$  (and  $c$  is finite), then  $\sum a_n$  and  $\sum b_n$  either both converge or both diverge.

Some series will “obviously” not converge—recognizing these can save you a lot of time and guesswork.

### Test for Divergence

If  $\lim_{n \rightarrow \infty} a_n \neq 0$ , then  $\sum_{n=1}^{\infty} a_n$  is divergent.

**Example:**  $\sum_{n=1}^{\infty} \frac{n^2 - 1}{n^2 + n}$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{n^2 - 1}{n^2 + n} = \lim_{n \rightarrow \infty} \frac{1 - \frac{1}{n^2}}{1 + \frac{1}{n}} = 1 \neq 0$$

so  $\sum_{n=1}^{\infty} \frac{n^2 - 1}{n^2 + n}$  is divergent.

## Testing for Convergence or Divergence of a Series (continued)

If  $a_n$  can be written as a function with a “nice” integral, the integral test may prove useful:

### Integral Test

If  $f(n) = a_n$  for all  $n$  and  $f(x)$  is continuous, positive, and decreasing on  $[1, \infty)$ , then:

If  $\int_1^{\infty} f(x) dx$  converges, then  $\sum_{n=1}^{\infty} a_n$  converges.

If  $\int_1^{\infty} f(x) dx$  is divergent, then  $\sum_{n=1}^{\infty} a_n$  is divergent.

**Example:**  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ .

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} \tan^{-1} x \Big|_1^t = \lim_{t \rightarrow \infty} \tan^{-1} t - \frac{\pi}{4} \\ &= \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}, \text{ so } \sum_{n=1}^{\infty} \frac{1}{n^2 + 1} \text{ is} \\ &\text{convergent.} \end{aligned}$$

**Example:**  $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^2}{n^3 + 1}$

$$\begin{aligned} \text{(i)} \quad \frac{1}{b_n} &= n + \frac{1}{n^2}, \text{ so } \frac{1}{b_{n+1}} = n + 1 + \frac{1}{(n+1)^2} > n + 1 \\ &\geq n + \frac{1}{n^2} = \frac{1}{b_n}, \text{ so } \frac{1}{b_{n+1}} \geq \frac{1}{b_n}, \text{ so } b_{n+1} \leq b_n \end{aligned}$$

$$\text{(ii)} \quad \lim_{n \rightarrow \infty} \frac{n^2}{n^3 + 1} = \lim_{n \rightarrow \infty} \frac{1}{n + 1/n^2} = 0$$

So  $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^2}{n^3 + 1}$  is convergent.

### Alternating Series Test

If (i)  $b_{n+1} \leq b_n$  for all  $n$  and (ii)

$\lim_{n \rightarrow \infty} b_n = 0$ , then  $\sum_{n=1}^{\infty} (-1)^{n-1} b_n$  is convergent.

The following 2 tests prove convergence, but also prove the stronger fact that  $\sum |a_n|$  converges (**absolute convergence**).

### Ratio Test

If  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$ , then  $\sum a_n$  is absolutely convergent.

If  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| > 1$  or  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$ , then  $\sum a_n$  is divergent.

If  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1$ , use another test.

**Example:**  $\sum_{n=1}^{\infty} e^{-n} n!$

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{e^{-n-1} (n+1)!}{e^{-n} n!} \right| \\ &= e^{-1} \lim_{n \rightarrow \infty} |n+1| = \infty, \text{ so} \\ &\sum_{n=1}^{\infty} e^{-n} n! \text{ is divergent.} \end{aligned}$$

**Example:**  $\sum_{n=1}^{\infty} \frac{n^n}{3^{1+3n}}$

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[n]{\frac{n^n}{3^{1+3n}}} &= \lim_{n \rightarrow \infty} \frac{n}{3^{1/n} 3^3} \\ &= \frac{1}{27} \lim_{n \rightarrow \infty} \frac{n}{3^{1/n}} = \infty \end{aligned}$$

So  $\sum_{n=1}^{\infty} \frac{n^n}{3^{1+3n}}$  is divergent.

When  $a_n$  contains factorials and/or powers of constants, as in the above example, the ratio test is often useful.

### Root Test

If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} < 1$ , then  $\sum a_n$  is absolutely convergent.

If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} > 1$  or  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$ , then  $\sum a_n$  is divergent.

If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = 1$ , use another test.